



LETTER TO THE EDITOR

Open Access

The significance of ^{18}F -FDG PET/CT in secondary hemophagocytic lymphohistiocytosis

Li-Juan Zhang^{1,2†}, Ji Xu^{1†}, Peng Liu¹, Chong-Yang Ding³, Jian-Yong Li¹, Hong-Xia Qiu^{1*} and Su-Jiang Zhang^{1*}

Abstract

This study was aimed to investigate the significance of ^{18}F -FDG PET/CT in secondary hemophagocytic lymphohistiocytosis (sHLH) patients. A total of 18 patients received ^{18}F -FDG PET/CT scan at initial diagnosis. All patients (18/18) had at least 3 organs involved, with increased FDG metabolism in different degrees. Fifteen cases (15/18) had definite underlying diseases, including infections (IAHLH), rheumatosis (RAHLH), or malignancy (MAHLH). The SUV_{max} of patients in MAHLH group was significantly higher than patients in IAHLH group or RAHLH group ($P=0.015$, $P=0.045$). Furthermore, the SUV_{max} of patients in IAHLH group was significantly higher than patients of RAHLH group ($P=0.043$). Therefore, we concluded that ^{18}F -FDG PET/CT may especially play important role in differential diagnosis of sHLH.

To the Editor

Secondary hemophagocytic lymphohistiocytosis (sHLH) is a hyper-inflammatory clinical syndrome mainly caused by severe infections, autoimmune inflammatory disorders and malignancies, especially lymphoma [1-3]. Up to date, very few data from the literature are available regarding the role of ^{18}F -FDG PET/CT in sHLH. In this study, 18 of 50 patients with sHLH who were admitted into our hospital between May 2007 and December 2010 underwent the examination (Table 1). The male-to-female ratio was 1:1, and the median age was 35 years (15-73). The diagnosis of HLH was made according to HLH-2004 diagnostic guidelines [4,5], and the underlying diseases were confirmed by a series of pathogenesis examinations including pathology, immunology, bacterial culture and virus detection et al. The maximum standardized uptake values (SUV_{max}) used to measure the level of FDG uptake were determined in all lesions [6]. All of the 18 patients had at least 3 organs involved, with increased FDG uptake at different level, including 18 cases showing splenomegaly, 16 cases serous effusions, 16 cases lymphadenopathy, 13 cases bone lesions, 12

cases pneumonia, 8 cases hepatomegaly, 5 cases brain parenchymal or cerebroventricular lesions, 5 cases cholecystitis, 4 cases myocardium lesions, and 2 cases kidney calculi. There were also other organs involved, such as larynx, muscles and adnexa. Fifteen patients (15/18) had definite underlying diseases, and were divided into three groups, including Infection Associated HLH (IAHLH, including EBV-HLH, $n=8$), Rheumatosis Associated HLH (RAHLH, $n=2$), and Malignancy Associated HLH (MAHLH, $n=5$). The SUV_{max} of patients in MAHLH group was significantly higher than those of patients with IAHLH (Mean 12.0 vs. 6.8, $P=0.015$), and RAHLH (Mean 12.0 vs. 2.7, $P=0.045$). Furthermore, the SUV_{max} of patients with IAHLH was significantly higher than that of patients with RAHLH (Mean 6.8 vs. 2.7, $P=0.043$). However, no significant difference in survival time was found between the three different sHLH subtype according to Kaplan-Meier analysis ($P>0.05$). In conclusion, ^{18}F -FDG PET/CT may play important role in differential diagnosis of sHLH, with high SUV pointing toward underlying malignancy.

* Correspondence: qhx9805@126.com; zbruce.zhang@gmail.com

†Equal contributors

¹Department of Hematology, The First Affiliated Hospital of Nanjing Medical University, Jiangsu Province Hospital, 300 Guangzhou Road, Nanjing, China
Full list of author information is available at the end of the article

Table 1 Characteristics of 18 sHLH patients

No.	Age/ Sex	Underlying disease	Therapy	Outcome	Survival (month)	Organs	SUV _{max}
1	35/M	Lymphoma (NK / T)	IVIG/HLH-2004 regimen(1 cycle) → High-dose methylprednisolone pulse therapy	Died of intracranial hemorrhage	1.7	6	12.3
2	35/F	Lymphoma (NK / T)	The Hyper-CVAD regimen (1 cycle)	Died of intracranial hemorrhage	1.2	6	15.7
3	18/M	Lymphoma (NK / T)	The CHOP regimen(1 cycle)	Died of acute hemorrhage of gastrointestinal tract	0.3	7	14.6
4	56/M	Lymphoma	Hydrocortisone 100mg×5d	Died of intracranial hemorrhage	1.7	5	4.3
5	32/M	Lymphoma	Dex 10mg/dx3d	Died of liver failure	0.3	10	13.3
6	37/F	Sjögren's syndrome	The COP regimen(3 cycle)	CR	>12	5	0.7
7	15/F	UCTD	The COP regimen (4 cycle)	CR	>45	3	4.6
8	21/F	EBV infection	HLH-2004 regimen (1 cycle)	Died of acute hemorrhage of gastrointestinal tract	1.7	7	6.6
9	17/M	EBV infection	Methylprednisolone 40 mg/dx24d	CR	>22	7	8.3
10	46/M	EBV infection	Dex 15mg/dx4d	Died of septic shock	0.4	6	10
11	73/M	EBV infection	The COP regimen (7 cycle)	Died of multi-organ failure	6	7	5.2
12	26/F	CMV infection	IVIG/HLH-2004 regimen (1 cycle) → The CHOP regimen(2 cycle)	CR	>24	6	9
13	24/F	CMV infection	The COP regimen (7 cycle)	Died of respiratory failure	2.2	5	4.2
14	69/F	MRSH infection	The COP regimen (2 cycle)	Died of respiratory failure	2.0	6	5.2
15	62/F	Fungal Infection	The COP regimen (7 cycle)	stable	>8	4	5.8
16	44/F	Malignant tumour?	Methylprednisolone 40 mg/dx5d	Died of multi-organ failure	0.4	8	7.7
17	56/M	Lymphoma?	The CHOP regimen (2 cycle) → Splenectomy→The Hyper-CVAD regimen (1cycle)	stable	>13	6	5.7
18	18/M	indefinite	HLH-2004 regimen (1 cycle)	Died of intracranial hemorrhage	0.2	3	4.2

HLH-2004, dexamethasone, etoposide and Cyclosporin A; CHOP, cyclophosphamide, doxorubicin, vincristine and prednisolone; COP, cyclophosphamide, vincristine and prednisone; Hyper-CVAD, cyclophosphamide, vincristine, doxorubicin, dexamethasone, methotrexate and cytarabine; DEX, dexamethasone; CR, complete response; UCTD, undifferentiated connective tissue disease; EBV, Epstein-Barr virus; CMV, cytomegalovirus; MRSH, methicillin-resistant *Staphylococcus hominis*.

Acknowledgments

This study was supported by the National Natural Science Foundation of China (81070456, 81170490), 'Liu Da Ren Cai Gao Feng' of Jiangsu Province (2010-WS-019), and A Project Funded by the Priority Academic Program Development of Jiangsu Higher Education Institutions.

Author details

¹Department of Hematology, The First Affiliated Hospital of Nanjing Medical University, Jiangsu Province Hospital, 300 Guangzhou Road, Nanjing, China. ²Department of Hematology, Huai'an First People's Hospital, Nanjing Medical University, Huai'an, China. ³Department of PET/CT Center, The First Affiliated Hospital of Nanjing Medical University, Jiangsu Province Hospital, Nanjing, China.

Received: 8 July 2012 Accepted: 23 July 2012
 Published: 23 July 2012

References

- Janka G, Imashuku S, Elinder G, *et al*: Infection- and malignancy-associated hemophagocytic syndromes. Secondary hemophagocytic lymphohistiocytosis. *Hematol Oncol Clin North Am* 1998, **12**:435–444.
- Fisman DN: Hemophagocytic syndromes and infection. *Emerg Infect Dis* 2000, **6**:601–608.
- Dhote R, Simon J, Papo T, *et al*: Reactive hemophagocytic syndrome in adult systemic disease: report of twenty-six cases and literature review. *Arthritis Rheum* 2003, **49**:633–639.

- Henter JL, Horne A, Aricó M, *et al*: HLH-2004: Diagnostic and therapeutic guidelines for hemophagocytic lymphohistiocytosis. *Pediatr Blood Cancer* 2007, **48**:124–131.
- Filipovich AH: Hemophagocytic lymphohistiocytosis (HLH) and related disorders. *Hematology Am Soc Hematol Educ Program* 2009, :127–131.
- Huang SC: Anatomy of SUV. *Nucl Med Biol* 2000, **27**:643–646.

doi:10.1186/1756-8722-5-40

Cite this article as: Zhang *et al*: The significance of ¹⁸F-FDG PET/CT in secondary hemophagocytic lymphohistiocytosis. *Journal of Hematology & Oncology* 2012 **5**:40.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

